



## TACC3 Polyclonal Antibody

Catalog_no :	AT4522
Applications :	WB,IHC-p,IF,ELISA
Reactivity :	Human,Mouse
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	TACC3
Protein_name :	Transforming acidic coiled-coil-containing protein 3
Humangene_id :	<a href="#">10460</a>
Humanswissprot_no :	<a href="#">Q9Y6A5</a>
Mouseswissprot_no :	<a href="#">Q9J111</a>
Immunogen :	The antiserum was produced against synthesized peptide derived from human TACC3. AA range:789-838
Specificity :	TACC3 Polyclonal Antibody detects endogenous levels of TACC3 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Rabbit
Dilution :	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
Msds :	<a href="#">MSDS_Antibody.pdf</a>
Other_name :	TACC3; ERIC1; Transforming acidic coiled-coil-containing protein 3; ERIC-1
Molecular Weight :	80KD


## Product Images



Western Blot analysis of various cells using TACC3 Polyclonal Antibody




Immunohistochemical analysis of paraffin-embedded Human testis. 1, Antibody was diluted at 1:100(4°, overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human testis. 1, Antibody was diluted at 1:100(4°, overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 30min).

Immunohistochemical analysis of paraffin-embedded Human testis. 1, Antibody was diluted at



1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).