



YB-1 Polyclonal Antibody

Catalog_no : AT4925

Applications : WB,IHC-p,ELISA

Reactivity : Human,Mouse,Rat

Category : 抗原抗体

Size : 100μg/50μg/20μg

Gene_name : YBX1

Protein_name : Nuclease-sensitive element-binding protein 1

Humangene_id : [4904](#)

:

Humanswissprot [P67809](#)
_no :

Mousegene_id : [22608](#)

Mouseswissprot [P62960](#)
_no :

Ratgene_id : [500538](#)

Ratswissprot_no [P62961](#)
:

Immunogen : The antiserum was produced against synthesized peptide derived from human YBX1. AA range:68-117

Specificity : YB-1 Polyclonal Antibody detects endogenous levels of YB-1 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source : Rabbit

Dilution : IHC-p: 100-300. Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.

Purification : The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage_stability : -20°C/1 year

:


Msds : MSDS_Antibody.pdf




Other_name : YBX1; NSEP1; YB1; Nuclease-sensitive element-binding protein 1; CCAAT-binding transcription factor I subunit A; CBF-A; DNA-binding protein B; DBPB; Enhancer factor I subunit A; EFI-A; Y-box transcription factor; Y-box-binding protein 1; YB-

Molecular Weight : 36KD


Product Images



Immunohistochemical analysis of paraffin-embedded Human testis. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human testis. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human testis. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).