

## ZNF785 Polyclonal Antibody

Catalog\_no: AT4984

Applications: WB,IHC-p,ELISA

Reactivity: Human

Category: 抗原抗体

Size:  $100 \mu g/50 \mu g/20 \mu g$ 

Gene\_name: ZNF785

Protein\_name: Zinc finger protein 785

Humangene\_id <u>146540</u>

:

Humanswissprot A8K8V0

\_no:

Immunogen: The antiserum was produced against synthesized peptide derived from human ZNF785.

AA range:91-140

Specificity: ZNF785 Polyclonal Antibody detects endogenous levels of ZNF785 protein.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Rabbit

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.

Not yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography

using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage\_stability -20°C/1 year

:

Msds: MSDS\_Antibody.pdf


Other\_name: ZNF785; Zinc finger protein 785

Molecular


87KD

Weight:

## **Product Images**



Western Blot analysis of HepG2 cells using ZNF785 Polyclonal Antibody. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.