

## CD66e Polyclonal Antibody

Catalog\_no: AT5172

Applications: WB,IHC-p,ELISA

Reactivity: Human

Category: 抗原抗体

Size: 100μg/50μg/20μg

Gene\_name: CEACAM5

Protein\_name: Carcinoembryonic antigen-related cell adhesion molecule 5

Humangene\_id 1048

Humanswissprot P06731

\_no:

The antiserum was produced against synthesized peptide derived from the Internal Immunogen:

region of human CEACAM5. AA range:481-530

Specificity: CD66e Polyclonal Antibody detects endogenous levels of CD66e protein.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Rabbit

Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other Dilution:

applications.

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography Purification:

using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage\_stability -20°C/1 year

Msds: MSDS\_Antibody.pdf

76KD

CEACAM5; CEA; Carcinoembryonic antigen-related cell adhesion molecule 5; Other\_name:


Carcinoembryonic antigen; CEA; Meconium antigen 100; CD66e

Molecular


Weight:

1/3


## **Product Images**



Western Blot analysis of MCF7 cells using CD66e Polyclonal Antibody.. Secondary antibody was diluted at 1:20000





Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100

Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100





Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100