

CXCR-3 Polyclonal Antibody


Catalog_no :	<u>AT5277</u>
Applications :	<u>WB,IHC-p,ELISA</u>
Reactivity :	<u>Human</u>
Category :	<u>抗原抗体</u>
Size :	<u>100µg/50µg/20µg</u>
Gene_name :	<u>CXCR3</u>
Protein_name :	<u>C-X-C chemokine receptor type 3</u>
Humangene_id :	<u>2833</u>
Humanswissprot_no :	<u>P49682</u>
Mousegene_id :	<u>12766</u>
Mouseswissprot_no :	<u>O88410</u>
Ratgene_id :	<u>84475</u>
Ratswissprot_no :	<u>Q9JII9</u>
Immunogen :	<u>The antiserum was produced against synthesized peptide derived from the Internal region of human CXCR3. AA range:161-210</u>
Specificity :	<u>CXCR-3 Polyclonal Antibody detects endogenous levels of CXCR-3 protein.</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.</u>
Purification :	<u>The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability :	<u>-20°C/1 year</u>
Msds :	<u>MSDS_Antibody.pdf</u>




Other_name : CXCR3; GPR9; C-X-C chemokine receptor type 3; CXC-R3; CXCR-3; CKR-L2; G protein-coupled receptor 9; Interferon-inducible protein 10 receptor; IP-10 receptor; CD183

Molecular Weight : 37KD


Product Images



Western Blot analysis of K562 cells using CXCR-3 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human brain, antibody was diluted at 1:100