

HNF-3 $\alpha$ / $\beta$ / $\gamma$  Polyclonal Antibody

Catalog_no :	<u>AT5324</u>
Applications :	<u>WB,ELISA</u>
Reactivity :	<u>Human,Mouse,Rat</u>
Category :	<u>抗原抗体</u>
Size :	<u>100<math>\mu</math>g/50<math>\mu</math>g/20<math>\mu</math>g</u>
Gene_name :	<u>FOXA1/FOXA2/FOXA3</u>
Protein_name :	<u>Hepatocyte nuclear factor 3-alpha/Hepatocyte nuclear factor 3-beta/Hepatocyte nuclear factor 3-gamma</u>
Humangene_id :	<u><a href="#">3169</a></u>
Humanswissprot_no :	<u><a href="#">P55317</a></u>
Mousegene_id :	<u><a href="#">15375</a></u>
Mouseswissprot_no :	<u><a href="#">P35582</a></u>
Ratgene_id :	<u><a href="#">25098</a></u>
Ratswissprot_no :	<u><a href="#">P23512</a></u>
Immunogen :	<u>The antiserum was produced against synthesized peptide derived from human FOXA1 around the non-acetylation site of Lys265. AA range:231-280</u>
Specificity :	<u>HNF-3<math>\alpha</math>/<math>\beta</math>/<math>\gamma</math> Polyclonal Antibody detects endogenous levels of HNF-3<math>\alpha</math>/<math>\beta</math>/<math>\gamma</math> protein.</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.</u>
Purification :	<u>The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability :	<u>-20°C/1 year</u>
Msds :	<u>MSDS_Antibody.pdf</u>




Other\_name : FOXA1; HNF3A; TCF3A; Hepatocyte nuclear factor 3-alpha; HNF-3-alpha; HNF-3A; Forkhead box protein A1; Transcription factor 3A; TCF-3A; FOXA2; HNF3B; TCF3B; Hepatocyte nuclear factor 3-beta; HNF-3-beta; HNF-3B; Forkhead box protein A2; Transcription factor 3B; TCF-3B; FOXA3; HNF3G; TCF3G; Hepatocyte nuclear factor 3-gamma; HNF-3-gamma; HNF-3G; Fork head-related protein FKH H3; Forkhead box protein A3; Transcription factor 3G; TCF-3G

---

Molecular Weight : 50KD

---

## Product Images



Western Blot analysis of A549 cells using HNF-3 $\alpha$ / $\beta$ / $\gamma$  Polyclonal Antibody.. Secondary antibody was diluted at 1:20000