

17 β -HSD4 Polyclonal Antibody


Catalog_no :	<u>AT5386</u>
Applications :	<u>WB,IHC-p,ELISA</u>
Reactivity :	<u>Human,Mouse,Rat</u>
Category :	<u>抗原抗体</u>
Size :	<u>100μg/50μg/20μg</u>
Gene_name :	<u>HSD17B4</u>
Protein_name :	<u>Peroxisomal multifunctional enzyme type 2</u>
Humangene_id :	<u>3295</u>
Humanswissprot_no :	<u>P51659</u>
Mousegene_id :	<u>15488</u>
Mouseswissprot_no :	<u>P51660</u>
Ratgene_id :	<u>79244</u>
Ratswissprot_no :	<u>P97852</u>
Immunogen :	<u>The antiserum was produced against synthesized peptide derived from the N-terminal region of human HSD17B4. AA range:41-90</u>
Specificity :	<u>17β-HSD4 Polyclonal Antibody detects endogenous levels of 17β-HSD4 protein.</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.</u>
Purification :	<u>The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability :	<u>-20°C/1 year</u>
Msds :	<u>MSDS_Antibody.pdf</u>




Other_name : HSD17B4; EDH17B4; Peroxisomal multifunctional enzyme type 2; MFE-2; 17-beta-hydroxysteroid dehydrogenase 4; 17-beta-HSD 4; D-bifunctional protein; DBP; Multifunctional protein 2; MPF-2

Molecular Weight : 80KD


Product Images



Western Blot analysis of HuvEc, K562 cells using 17 β -HSD4 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100

Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100

