

CD298 Polyclonal Antibody


Catalog_no :	<u>AT5623</u>
Applications :	<u>WB,ELISA</u>
Reactivity :	<u>Human</u>
Category :	<u>抗原抗体</u>
Size :	<u>100µg/50µg/20µg</u>
Gene_name :	<u>ATP1B3</u>
Protein_name :	<u>Sodium/potassium-transporting ATPase subunit beta-3</u>
Humangene_id	<u>483</u>
:	<u></u>
Humanswissprot	<u>P54709</u>
_no :	<u></u>
Mousegene_id :	<u>11933</u>
Mouseswissprot	<u>P97370</u>
_no :	<u></u>
Ratgene_id :	<u>25390</u>
Ratswissprot_no	<u>Q63377</u>
:	<u></u>
Immunogen :	<u>The antiserum was produced against synthesized peptide derived from the C-terminal region of human ATP1B3. AA range:222-271</u>
Specificity :	<u>CD298 Polyclonal Antibody detects endogenous levels of CD298 protein.</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.</u>
Purification :	<u>The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability	<u>-20°C/1 year</u>
:	<u></u>
Msds :	<u>MSDS_Antibody.pdf</u>



Other_name : ATP1B3; Sodium/potassium-transporting ATPase subunit beta-3; Sodium/potassium-dependent ATPase subunit beta-3; ATPB-3; CD298

Molecular Weight : 36KD

Product Images



Western Blot analysis of PC12, NIH-3T3 cells using CD298 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000