

HGF Polyclonal Antibody

Catalog_no:	AT5699
Applications :	WB,IHC-p,ELISA
Reactivity :	Human
Category :	抗原抗体
Size :	100μg/50μg/20μg
Gene_name :	HGF
Protein_name :	Hepatocyte growth factor
Humangene_id :	<u>3082</u>
Humanswisspro _no:	t <u>P14210</u>
Mousegene_id:	
Mouseswissprot _no:	<u>Q08048</u>
Ratgene_id:	
Ratswissprot_no :	
(mmunogen :	The antiserum was produced against synthesized peptide derived from the Internal region of human HGF. AA range:500-600
Specificity :	HGF Polyclonal Antibody detects endogenous levels of HGF protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Rabbit
Dilution :	IHC-p: 100-300.Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
VIsds:	MSDS_Antibody.pdf
	<u> </u>




Other_name: HGF; HPTA; Hepatocyte growth factor; Hepatopoietin-A; Scatter factor; SF


Molecular Weight: 80KD




Product Images



Western blot analysis of SH-SY5Y lysis using HGF antibody. Antibody was diluted at 1:500. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).





Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).