

ICOS Polyclonal Antibody


Catalog_no :	AT5713
Applications :	WB,IHC-p,ELISA
Reactivity :	Human,Mouse,Rat
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	ICOS
Protein_name :	Inducible T-cell costimulator
Humangene_id :	29851
Humanswissprot_no :	Q9Y6W8
Mousegene_id :	54167
Mouseswissprot_no :	Q9WVS0
Ratgene_id :	64545
Ratswissprot_no :	Q9R1T7
Immunogen :	The antiserum was produced against synthesized peptide derived from the Internal region of human ICOS. AA range:31-80
Specificity :	ICOS Polyclonal Antibody detects endogenous levels of ICOS protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Rabbit
Dilution :	Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
Msds :	MSDS_Antibody.pdf




Other_name : ICOS; AILIM; Inducible T-cell costimulator; Activation-inducible lymphocyte immunomediatory molecule; CD278

Molecular Weight : 22KD

Product Images



Western blot analysis of 3T3 lysis using ICOS antibody. Antibody was diluted at 1:500. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human tonsils, antibody was diluted at 1:100