

CTACK Polyclonal Antibody


Catalog_no :	AT5724
Applications :	WB,IHC-p,ELISA
Reactivity :	Human
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	CCL27
Protein_name :	C-C motif chemokine 27
Humangene_id :	10850
Humanswissprot_no :	Q9Y4X3
Mousegene_id :	100039863
Mouseswissprot_no :	Q9Z1X0
Ratgene_id :	
Ratswissprot_no :	
Immunogen :	Synthesized peptide derived from C-C motif chemokine 27 at AA range: 51-100
Specificity :	CTACK Polyclonal Antibody detects endogenous levels of CTACK protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Rabbit
Dilution :	Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
Msds :	MSDS_Antibody.pdf




Other_name : CCL27; ILC; SCYA27; C-C motif chemokine 27; CC chemokine ILC; Cutaneous T-cell-attracting chemokine; CTACK; ESkin; IL-11 R-alpha-locus chemokine; Skinkine; Small-inducible cytokine A27

Molecular Weight : 12KD

Product Images



Western blot analysis of mouse-kidney lysis using CCL27 antibody. Antibody was diluted at 1:1000. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-skin, antibody was diluted at 1:100

Western Blot analysis of various cells using Antibody diluted at 1:1000. Secondary antibody was diluted at 1:20000

