





## Endoglin Polyclonal Antibody

|                     |  |
|---------------------|--|
| Catalog_no :        | <u>AT5879</u>  |
| Applications :      | <u>WB,IHC-p,ELISA</u>  |
| Reactivity :        | <u>Human</u>   |
| Category :          | <u>抗原抗体</u>  |
| Size :              | <u>100µg/50µg/20µg</u>   |
| Gene_name :         | <u>ENG END</u>   |
| Protein_name :      | <u>Endoglin (CD antigen CD105)</u>   |
| Humangene_id :      | <u><a href="#">2022</a></u>  |
| Humanswissprot_no : | <u><a href="#">P17813</a></u>  |
| Immunogen :         | <u>Synthetic peptide from human protein at AA range: 370-430</u>   |
| Specificity :       | <u>The antibody detects endogenous Endoglin</u>  |
| Formulation :       | <u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>   |
| Source :            | <u>Rabbit</u>  |
| Dilution :          | <u>WB 1:500-2000,IHC-p 1:500-200, ELISA 1:10000-20000</u>  |
| Purification :      | <u>The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.</u> |
| Concentration :     | <u>1 mg/ml</u>   |
| Storage_stability : | <u>-20°C/1 year</u>  |
| Msds :              | <u>MSDS_Antibody.pdf</u>   |
| Other_name :        | <u>Endoglin (CD antigen CD105)</u>   |
| Molecular Weight :  | <u>70KD</u>  |


## Product Images



Western blot analysis of 3T3 KB K562 Hela 293T lysate, antibody was diluted at 500. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-tonsil, antibody was diluted at 1:200




Immunohistochemical analysis of paraffin-embedded human-tonsil, antibody was diluted at 1:200

Immunohistochemical analysis of paraffin-embedded






human-kidney, antibody was diluted at 1:200






Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).





Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:200(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

