



## MIG Polyclonal Antibody

Catalog\_no : AT6008

Applications : IHC-p,ELISA

Reactivity : Human

Category : 抗原抗体

Size : 100μg/50μg/20μg

Gene\_name : CXCL9 CMK MIG SCYB9

Protein\_name : C-X-C motif chemokine 9 (Gamma-interferon-induced monokine) (Monokine induced by interferon-gamma) (HuMIG) (MIG) (Small-inducible cytokine B9)

Humangene\_id : [4283](#)

:

Humanswissprot [Q07325](#)

\_no :

Immunogen : Synthetic peptide from human protein at AA range: 90-125

Specificity : The antibody detects endogenous MIG

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source : Rabbit

Dilution : IHC-p 1:50-200, ELISA 1:10000-20000

Purification : The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml


Storage\_stability : -20°C/1 year

:


Msds : [MSDS\\_Antibody.pdf](#)

Other\_name : C-X-C motif chemokine 9 (Gamma-interferon-induced monokine) (Monokine induced by interferon-gamma) (HuMIG) (MIG) (Small-inducible cytokine B9)

## Product Images



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200