


CD56 (3C8) mouse mAb

| | |
|---------------------|--|
| Catalog_no : | AM6574 |
| Applications : | IHC-p,WB |
| Reactivity : | Human |
| Category : | 抗原抗体 |
| Size : | 50µl/100µl |
| Gene_name : | NCAM1 NCAM |
| Protein_name : | Neural cell adhesion molecule 1 (N-CAM-1) (NCAM-1) (CD antigen CD56) |
| Humanswissprot_no : | P13591 |
| Immunogen : | Synthesized peptide derived from human CD56 |
| Specificity : | This antibody detects endogenous levels of human CD56 |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Mouse IgG |
| Dilution : | IHC-p 1:100-500 , WB 1:500-2000 |
| Purification : | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability : | -20°C/1 year |
| Notes : | cell adhesion, cell surface receptor linked signal transduction, positive regulation of signal transduction, positive regulation of cell communication, cell-cell adhesion, biological adhesion, homotypic cell-cell adhesion, regulation of calcium-media |

Product Images



Immunohistochemical analysis of paraffin-embedded Small cell lung carcinoma. 1, Antibody was diluted at 1:200(4°,overnight). 2, TRIS-EDTA of pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).