

## CTBL1 rabbit pAb

Catalog\_no: AN3149

WB Applications:

Reactivity: Human, Mouse, Rat

Category: 抗原抗体

Size: 100μg/50μg/20μg

CTNNBL1 C20orf33 PP8304 Gene\_name:

Protein\_name: CTBL1

Humangene\_id 56259

Humanswissprot Q8WYA6

\_no:

Mousegene\_id: 66642

Mouseswissprot **Q9CWL8** 

\_no:

Ratgene\_id: 296320

Ratswissprot\_no Q4V8K2

Immunogen: Synthesized peptide derived from human CTBL1

This antibody detects endogenous levels of CTBL1 at Human/Mouse/Rat Specificity:

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.107% sodium azide.

Source: Rabbit

Dilution: WB 1:500-2000

Purification: The antibody was affinity-purified from rabbit serum by affinity-chromatography using

specific immunogen.

Concentration: 1 mg/ml

Storage\_stability -20°C/1 year

Other\_name: Beta-catenin-like protein 1 (Nuclear-associated protein) (NAP) (Testis development

protein NYD-SP19)



Molecular Weight: 60KD

Background:

The protein encoded by this gene is a component of the pre-mRNA-processing factor 19-cell division cycle 5-like (PRP19-CDC5L) protein complex, which activates pre-mRNA splicing and is an integral part of the spliceosome. The encoded protein is also a nuclear localization sequence binding protein, and binds to activation-induced deaminase and is important for antibody diversification. This gene may also be associated with the development of obesity. Alternative splicing results in multiple transcript variants. A pseudogene of this gene has been defined on the X chromosome. [provided by RefSeq, Jul 2013],