



OSBL1 rabbit pAb

Catalog_no :	<u>AN3251</u>
Applications :	<u>WB</u>
Reactivity :	<u>Human, Mouse,Rat</u>
Category :	<u>抗原抗体</u>
Size :	<u>100µg/50µg/20µg</u>
Gene_name :	<u>OSBPL1A ORP1 OSBP8 OSBPL1 OSBPL1B</u>
Protein_name :	<u>OSBL1</u>
Humangene_id :	<u>114876</u>
Humanswissprot_no :	<u>Q9BXW6</u>
Mousegene_id :	<u>64291</u>
Mouseswissprot_no :	<u>Q91XL9</u>
Ratgene_id :	<u>259221</u>
Ratswissprot_no :	<u>Q8K4M9</u>
Immunogen :	<u>Synthesized peptide derived from human OSBL1</u>
Specificity :	<u>This antibody detects endogenous levels of OSBL1 at Human/Mouse/Rat</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.209% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>WB 1:500-2000</u>
Purification :	<u>The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability :	<u>-20°C/1 year</u>
Other_name :	<u>Oxysterol-binding protein-related protein 1 (ORP-1) (OSBP-related protein 1)</u>
Molecular	<u>105KD</u>



Weight :

Background :

This gene encodes a member of the oxysterol-binding protein (OSBP) family, a group of intracellular lipid receptors. Most members contain an N-terminal pleckstrin homology domain and a highly conserved C-terminal OSBP-like sterol-binding domain, although some members contain only the sterol-binding domain. Transcript variants derived from alternative promoter usage and/or alternative splicing exist; they encode different isoforms. [provided by RefSeq, Jul 2008],
