

## WDR36 rabbit pAb

Catalog\_no: AN3261

Applications: WB

Reactivity: Human

Category: 抗原抗体

Size: 100μg/50μg/20μg

Gene\_name: WDR36

Protein\_name: WDR36

Humangene\_id 134430

Humanswissprot **Q8NI36** 

\_no:

Synthesized peptide derived from human WDR36 Immunogen:

Specificity: This antibody detects endogenous levels of WDR36 at Human

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.219% sodium azide.

Source: Rabbit

Dilution: WB 1:500-2000

Purification: The antibody was affinity-purified from rabbit serum by affinity-chromatography using

specific immunogen.

Concentration: 1 mg/ml

Storage\_stability -20°C/1 year

WD repeat-containing protein 36 (T-cell activation WD repeat-containing protein) (TA-Other\_name:

WDRP)

Molecular Weight:

105KD

Background: This gene encodes a member of the WD repeat protein family. WD repeats are

minimally conserved regions of approximately 40 amino acids typically bracketed by gly-

regulation. Mutations in this gene have been associated with adult-onset primary open-

his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene

angle glaucoma (POAG). [provided by RefSeq, Jul 2008],

