

PCDC1 rabbit pAb

Catalog_no :	AN3309
Applications :	WB
Reactivity :	Human
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	PCDHAC1
Protein_name :	PCDC1
Humangene_id :	56135
Humanswissprot_no :	Q9H158
Immunogen :	Synthesized peptide derived from human PCDC1
Specificity :	This antibody detects endogenous levels of PCDC1 at Human
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.267% sodium azide.
Source :	Rabbit
Dilution :	WB 1:500-2000
Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
Other_name :	Protocadherin alpha-C1 (PCDH-alpha-C1)
Molecular Weight :	105KD
Background :	This gene is a member of the protocadherin alpha gene cluster, one of three related gene clusters tandemly linked on chromosome five that demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The alpha gene cluster is composed of 15 cadherin superfamily genes related to the mouse CNR genes and consists of 13 highly similar and 2 more distantly related coding sequences. The tandem array of 15 N-terminal exons, or variable exons, are followed by downstream C-terminal exons, or constant exons, which are shared by all genes in the cluster. The large, uninterrupted N-terminal exons each encode six cadherin

ectodomains while the C-terminal exons encode the cytoplasmic domain. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins that most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been observed and additional variants have been suggested but their full-length nature has yet to be determined. [provided by RefSeq, Jul 2008],
