

## NAA25 rabbit pAb

Catalog_no :	AN3331
Applications :	WB
Reactivity :	Human, Mouse,Rat
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	NAA25 C12orf30 MDM20 NAP1
Protein_name :	NAA25
Humangene_id	<a href="#">80018</a>
:	
Humanswissprot	<a href="#">Q14CX7</a>
_no :	
Mousegene_id :	<a href="#">231713</a>
Mouseswissprot	<a href="#">Q8BWZ3</a>
_no :	
Ratgene_id :	<a href="#">360811</a>
Ratswissprot_no	<a href="#">Q6QI44</a>
:	
Immunogen :	Synthesized peptide derived from human NAA25
Specificity :	This antibody detects endogenous levels of NAA25 at Human/Mouse/Rat
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.289% sodium azide.
Source :	Rabbit
Dilution :	WB 1:500-2000
Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Concentration :	1 mg/ml
Storage_stability	-20°C/1 year
:	
Other_name :	N-alpha-acetyltransferase 25, NatB auxiliary subunit (Mitochondrial distribution and morphology protein 20) (N-terminal acetyltransferase B complex subunit MDM20) (NatB complex subunit MDM20) (N-terminal acetyltransferase B complex subunit NAA25)



(p120)

---

Molecular Weight : 105KD

---

Background : This gene encodes the auxiliary subunit of the heteromeric N-terminal acetyltransferase B complex. This complex acetylates methionine residues that are followed by acidic or asparagine residues.[provided by RefSeq, Mar 2010],

---