

UBC rabbit pAb

Catalog_no :	<u>AN3742</u>
Applications :	<u>WB</u>
Reactivity :	<u>Human, Mouse,Rat</u>
Category :	<u>抗原抗体</u>
Size :	<u>100µg/50µg/20µg</u>
Gene_name :	<u>UBC</u>
Protein_name :	<u>UBC</u>
Humangene_id :	<u>0</u>
Humanswissprot_no :	<u>P0CG48</u>
Mousegene_id :	<u>22190</u>
Mouseswissprot_no :	<u>P0CG50</u>
Ratgene_id :	<u>50522</u>
Ratswissprot_no :	<u>Q63429</u>
Immunogen :	<u>Synthesized peptide derived from human UBC</u>
Specificity :	<u>This antibody detects endogenous levels of UBC at Human/Mouse/Rat</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>WB 1 : 500-2000</u>
Purification :	<u>The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability :	<u>-20°C/1 year</u>
Background :	<u>This gene represents a ubiquitin gene, ubiquitin C. The encoded protein is a polyubiquitin precursor. Conjugation of ubiquitin monomers or polymers can lead to various effects within a cell, depending on the residues to which ubiquitin is conjugated.</u>



Ubiquitination has been associated with protein degradation, DNA repair, cell cycle regulation, kinase modification, endocytosis, and regulation of other cell signaling pathways. [provided by RefSeq, Aug 2010],
