

BYST rabbit pAb

Catalog_no :	AN3743
Applications :	WB
Reactivity :	Human, Mouse,Rat
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	BYSL
Protein_name :	BYST
Humangene_id :	705
Humanswissprot_no :	Q13895
Mousegene_id :	53414
Mouseswissprot_no :	O54825
Ratgene_id :	359727
Ratswissprot_no :	Q80WL2
Immunogen :	Synthesized peptide derived from human BYST
Specificity :	This antibody detects endogenous levels of BYST at Human/Mouse/Rat
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Rabbit
Dilution :	WB 1 : 500-2000
Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
Background :	Bystin is expressed as a 2-kb major transcript and a 3.6-kb minor transcript in SNG-M cells and in human trophoblastic teratocarcinoma HT-H cells. Protein binding assays determined that bystin binds directly to trophinin and tastin, and that binding is

enhanced when cytokeratins 8 and 18 are present. Immunocytochemistry of HT-H cells showed that bystin colocalizes with trophinin, tastin, and the cytokeratins, suggesting that these molecules form a complex in trophoblast cells at the time of implantation. Using immunohistochemistry it was determined that trophinin and bystin are found in the placenta from the sixth week of pregnancy. Both proteins were localized in the cytoplasm of the syncytiotrophoblast in the chorionic villi and in endometrial decidual cells at the uteroplacental interface. After week 10, the levels of trophinin, tastin, and bystin decreased and then disappeared from placental villi. [provided by RefSeq, Jul 2008],
