

AP2B rabbit pAb

Catalog_no :	<u>AN3852</u>
Applications :	<u>WB</u>
Reactivity :	<u>Human, Mouse</u>
Category :	<u>抗原抗体</u>
Size :	<u>100µg/50µg/20µg</u>
Gene_name :	<u>TFAP2B</u>
Protein_name :	<u>AP2B</u>
Humangene_id :	<u>7021</u>
Humanswissprot_no :	<u>Q92481</u>
Mousegene_id :	<u>21419</u>
Mouseswissprot_no :	<u>Q61313</u>
Immunogen :	<u>Synthesized peptide derived from human AP2B</u>
Specificity :	<u>This antibody detects endogenous levels of AP2B at Human/Mouse</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>WB 1 : 500-2000</u>
Purification :	<u>The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability :	<u>-20°C/1 year</u>
Background :	<p>This gene encodes a member of the AP-2 family of transcription factors. AP-2 proteins form homo- or hetero-dimers with other AP-2 family members and bind specific DNA sequences. They are thought to stimulate cell proliferation and suppress terminal differentiation of specific cell types during embryonic development. Specific AP-2 family members differ in their expression patterns and binding affinity for different promoters. This protein functions as both a transcriptional activator and repressor. Mutations in this gene result in autosomal dominant Char syndrome, suggesting that this gene functions in the differentiation of neural crest cell derivatives. [provided by RefSeq, Jul</p>



2008],
