

PCDG9 rabbit pAb

Catalog_no: AN3880

Applications: WB

Reactivity: Human

Category: 抗原抗体

Size: $100 \mu g/50 \mu g/20 \mu g$

Gene_name : PCDHGA9

Protein_name: PCDG9

Humangene_id 56107

:

Humanswissprot <u>Q9Y5G4</u>

no:

Immunogen: Synthesized peptide derived from human PCDG9

Specificity: This antibody detects endogenous levels of PCDG9 at Human

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Rabbit

Dilution: WB 1:500-2000

Purification: The antibody was affinity-purified from rabbit serum by affinity-chromatography using

specific immunogen.

Concentration: 1 mg/ml

Storage_stability -20°C/1 year

:

Background: This gene is a member of the protocadherin gamma gene cluster, one of three related

clusters tandemly linked on chromosome five. These gene clusters have an immunoglobulin-like organization, suggesting that a novel mechanism may be involved in their regulation and expression. The gamma gene cluster includes 22 genes divided into 3 subfamilies. Subfamily A contains 12 genes, subfamily B contains 7 genes and 2 pseudogenes, and the more distantly related subfamily C contains 3 genes. The tandem array of 22 large, variable region exons are followed by a constant region, containing 3 exons shared by all genes in the cluster. Each variable region exon encodes the extracellular region, which includes 6 cadherin ectodomains and a transmembrane region. The constant region exons encode the common cytoplasmic region. These neural cadherin-like cell adhesion proteins most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been described for the gamma cluster genes. [provided by RefSeq, Jul 2008],

