

## ASCC1 rabbit pAb

Catalog_no :	AN4067
Applications :	WB
Reactivity :	Human, Mouse
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	ASCC1 CGI-18
Protein_name :	ASCC1
Humangene_id :	<a href="#">51008</a>
Humanswissprot_no :	<a href="#">Q8N9N2</a>
Mousegene_id :	<a href="#">69090</a>
Mouseswissprot_no :	<a href="#">Q9D8Z1</a>
Immunogen :	Synthesized peptide derived from human ASCC1
Specificity :	This antibody detects endogenous levels of ASCC1 at Human/Mouse
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Rabbit
Dilution :	WB 1 : 500-2000
Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
Background :	This gene encodes a subunit of the activating signal cointegrator 1 (ASC-1) complex. The ASC-1 complex is a transcriptional coactivator that plays an important role in gene transactivation by multiple transcription factors including activating protein 1 (AP-1), nuclear factor kappa-B (NF-kB) and serum response factor (SRF). The encoded protein contains an N-terminal KH-type RNA-binding motif which is required for AP-1 transactivation by the ASC-1 complex. Mutations in this gene are associated with Barrett esophagus and esophageal adenocarcinoma. Alternatively spliced transcripts encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2011],



**专注二抗产品**

Biodragon旗下品牌

---