



## BTBD1 rabbit pAb

Catalog_no :	<u>AN4090</u>
Applications :	<u>WB</u>
Reactivity :	<u>Human, Mouse</u>
Category :	<u>抗原抗体</u>
Size :	<u>100µg/50µg/20µg</u>
Gene_name :	<u>BTBD1 C15orf1 NS5ATP8</u>
Protein_name :	<u>BTBD1</u>
Humangene_id :	<u><a href="#">53339</a></u>
Humanswissprot_no :	<u><a href="#">Q9H0C5</a></u>
Mousegene_id :	<u><a href="#">83962</a></u>
Mouseswissprot_no :	<u><a href="#">P58544</a></u>
Immunogen :	<u>Synthesized peptide derived from human BTBD1</u>
Specificity :	<u>This antibody detects endogenous levels of BTBD1 at Human/Mouse</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>WB 1 : 500-2000</u>
Purification :	<u>The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability :	<u>-20°C/1 year</u>
Background :	<u>The C-terminus of the protein encoded by this gene binds topoisomerase I. The N-terminus contains a proline-rich region and a BTB/POZ domain (broad-complex, Tramtrack and bric a brac/Pox virus and Zinc finger), both of which are typically involved in protein-protein interactions. Subcellularly, the protein localizes to cytoplasmic bodies. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008],</u>



**专注二抗产品**

Biodragon旗下品牌

---