

## NMD3 rabbit pAb

|                     |   |
|---------------------|---|
| Catalog_no :        | AN4256  |
| Applications :      | WB  |
| Reactivity :        | Human, Mouse  |
| Category :          | 抗原抗体  |
| Size :              | 100µg/50µg/20µg   |
| Gene_name :         | NMD3 CGI-07   |
| Protein_name :      | NMD3  |
| Humangene_id :      | <a href="#">51068</a>   |
| Humanswissprot_no : | <a href="#">Q96D46</a>  |
| Mousegene_id :      | <a href="#">97112</a>   |
| Mouseswissprot_no : | <a href="#">Q99L48</a>  |
| Immunogen :         | Synthesized peptide derived from human NMD3   |
| Specificity :       | This antibody detects endogenous levels of NMD3 at Human/Mouse  |
| Formulation :       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source :            | Rabbit  |
| Dilution :          | WB 1 : 500-2000   |
| Purification :      | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.   |
| Concentration :     | 1 mg/ml   |
| Storage_stability : | -20°C/1 year  |
| Background :        | Ribosomal 40S and 60S subunits associate in the nucleolus and are exported to the cytoplasm. The protein encoded by this gene is involved in the passage of the 60S subunit through the nuclear pore complex and into the cytoplasm. Several transcript variants exist for this gene, but the full-length nature of only two have been described to date. [provided by RefSeq, Feb 2016], |