

## LIS1 rabbit pAb

Catalog\_no: AT6353

WB Applications:

Reactivity: Human, Mouse, Rat

Category: 抗原抗体

Size: 100μg/50μg/20μg

PAFAH1B1 LIS1 MDCR MDS PAFAHA Gene\_name:

Protein\_name: LIS1

Humangene\_id 5048

Humanswissprot P43034

\_no:

Mousegene\_id: 18472

Mouseswissprot P63005

\_no:

Ratgene\_id: 83572

Ratswissprot\_no P63004

Immunogen: Synthesized peptide derived from human LIS1

This antibody detects endogenous levels of LIS1 at Human/Mouse/Rat Specificity:

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Rabbit

Dilution: WB 1:500-2000

The antibody was affinity-purified from rabbit serum by affinity-chromatography using Purification:

specific immunogen.

Concentration: 1 mg/ml

Storage\_stability -20°C/1 year

Background: This locus was identified as encoding a gene that when mutated or lost caused the

> lissencephaly associated with Miller-Dieker lissencephaly syndrome. This gene encodes the non-catalytic alpha subunit of the intracellular Ib isoform of platelet-activating factor



acteylhydrolase, a heterotrimeric enzyme that specifically catalyzes the removal of the acetyl group at the SN-2 position of platelet-activating factor (identified as 1-O-alkyl-2-acetyl-sn-glyceryl-3-phosphorylcholine). Two other isoforms of intracellular platelet-activating factor acetylhydrolase exist: one composed of multiple subunits, the other, a single subunit. In addition, a single-subunit isoform of this enzyme is found in serum. [provided by RefSeq, Apr 2009],