

## RCBT2 rabbit pAb

Catalog_no :	<u>AT6715</u>
Applications :	<u>WB</u>
Reactivity :	<u>Human, Mouse,Rat</u>
Category :	<u>抗原抗体</u>
Size :	<u>100µg/50µg/20µg</u>
Gene_name :	<u>RCBTB2 CHC1L RLG</u>
Protein_name :	<u>RCBT2</u>
Humangene_id	<u><a href="#">1102</a></u>
:	
Humanswissprot	<u><a href="#">O95199</a></u>
_no :	
Mousegene_id :	<u><a href="#">105670</a></u>
Mouseswissprot	<u><a href="#">Q99LJ7</a></u>
_no :	
Ratgene_id :	<u><a href="#">290363</a></u>
Ratswissprot_no	<u><a href="#">Q6P798</a></u>
:	
Immunogen :	<u>Synthesized peptide derived from human RCBT2</u>
Specificity :	<u>This antibody detects endogenous levels of RCBT2 at Human/Mouse/Rat</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>WB 1 : 500-2000</u>
Purification :	<u>The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability	<u>-20°C/1 year</u>
:	
Background :	<u>This gene encodes a protein containing two C-terminal BTB/POZ domains that is related to regulator of chromosome condensation (RCC). The encoded protein may act as a guanine nucleotide exchange factor. This gene is observed to be lost or underexpressed</u>



in prostate cancers. There is a pseudogene of this gene on chromosome 10. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2013],