

## S38A8 rabbit pAb

|                     |  |
|---------------------|--|
| Catalog_no :        | AT6725   |
| Applications :      | WB   |
| Reactivity :        | Human, Mouse   |
| Category :          | 抗原抗体   |
| Size :              | 100µg/50µg/20µg  |
| Gene_name :         | SLC38A8  |
| Protein_name :      | S38A8  |
| Humangene_id :      | <a href="#">146167</a>   |
| Humanswissprot_no : | <a href="#">A6NNN8</a>   |
| Mousegene_id :      | <a href="#">234788</a>   |
| Mouseswissprot_no : | <a href="#">Q5HZH7</a>   |
| Immunogen :         | Synthesized peptide derived from human S38A8   |
| Specificity :       | This antibody detects endogenous levels of S38A8 at Human/Mouse  |
| Formulation :       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source :            | Rabbit   |
| Dilution :          | WB 1 : 500-2000  |
| Purification :      | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  |
| Concentration :     | 1 mg/ml  |
| Storage_stability : | -20°C/1 year   |
| Background :        | This gene encodes a putative sodium-dependent amino-acid/proton antiporter. The protein has eleven transmembrane domains, an extracellular N-terminus and an intracellular C-terminal tail. The protein is a member of the SLC38 sodium-coupled neutral amino acid transporter family of proteins. Mutations in this gene result in foveal hypoplasia with or without optic nerve misrouting and/or anterior segment dysgenesis. [provided by RefSeq, May 2014], |



**专注二抗产品**

Biodragon旗下品牌

---