

FBX6 rabbit pAb

| | |
|---------------------|---|
| Catalog_no : | AT6731 |
| Applications : | WB |
| Reactivity : | Human, Mouse,Rat |
| Category : | 抗原抗体 |
| Size : | 100µg/50µg/20µg |
| Gene_name : | FBXO6 FBG2 FBS2 FBX6 |
| Protein_name : | FBX6 |
| Humangene_id : | 26270 |
| Humanswissprot_no : | Q9NRD1 |
| Mousegene_id : | 50762 |
| Mouseswissprot_no : | Q9QZN4 |
| Ratgene_id : | 192351 |
| Ratswissprot_no : | Q923V4 |
| Immunogen : | Synthesized peptide derived from human FBX6 |
| Specificity : | This antibody detects endogenous levels of FBX6 at Human/Mouse/Rat |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Rabbit |
| Dilution : | WB 1 : 500-2000 |
| Purification : | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability : | -20°C/1 year |
| Background : | This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), |

which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class, and its C-terminal region is highly similar to that of rat NFB42 (neural F Box 42 kDa) which may be involved in the control of the cell cycle. [provided by RefSeq, Jul 2008],
