

CHD7 rabbit pAb

| | |
|---------------------|--|
| Catalog_no : | <u>AT7184</u> |
| Applications : | <u>IHC-p</u> |
| Reactivity : | <u>Human, Mouse</u> |
| Category : | <u>抗原抗体</u> |
| Size : | <u>100µg/50µg/20µg</u> |
| Gene_name : | <u>CHD7 KIAA1416</u> |
| Protein_name : | <u>CHD7</u> |
| Humangene_id : | <u>55636</u> |
| Humanswissprot_no : | <u>Q9P2D1</u> |
| Mousegene_id : | <u>320790</u> |
| Mouseswissprot_no : | <u>A2AJK6</u> |
| Immunogen : | <u>Synthesized peptide derived from human CHD7</u> |
| Specificity : | <u>This antibody detects endogenous levels of CHD7 at Human/Mouse</u> |
| Formulation : | <u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u> |
| Source : | <u>Rabbit</u> |
| Dilution : | <u>IHC-p 1 : 50-200</u> |
| Purification : | <u>The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.</u> |
| Concentration : | <u>1 mg/ml</u> |
| Storage_stability : | <u>-20°C/1 year</u> |
| Background : | <u>This gene encodes a protein that contains several helicase family domains. Mutations in this gene have been found in some patients with the CHARGE syndrome. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2015],</u> |