

## PCSK5 rabbit pAb

Catalog\_no: AT7200

WB Applications:

Reactivity: Human, Mouse, Rat

Category: 抗原抗体

100μg/50μg/20μg Size:

PCSK5 PC5 PC6 Gene\_name:

Protein\_name : PCSK5

Humangene\_id 5125

Humanswissprot Q92824

\_no:

Mousegene\_id: 18552

Mouseswissprot **Q04592** 

\_no:

Ratgene\_id: 116548

Ratswissprot\_no P41413

Immunogen: Synthesized peptide derived from human PCSK5

This antibody detects endogenous levels of PCSK5 at Human/Mouse/Rat Specificity:

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Rabbit

Dilution: WB 1:500-2000

The antibody was affinity-purified from rabbit serum by affinity-chromatography using Purification:

specific immunogen.

Concentration: 1 mg/ml

Storage\_stability -20°C/1 year

Background: This gene encodes a member of the subtilisin-like proprotein convertase family, which

> includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein



undergoes an initial autocatalytic processing event in the ER to generate a heterodimer which exits the ER. It then sorts to the trans-Golgi network where a second autocatalytic event takes place and the catalytic activity is acquired. This encoded protein is widely expressed and one of the seven basic amino acid-specific members which cleave their substrates at single or paired basic residues. It mediates posttranslational endoproteolytic processing for several integrin alpha subunits and is thought to process prorenin, pro-membrane type-1 matrix metalloproteinase and HIV-1 glycoprotein gp160. Alternative splicing results in multiple transcript variants, some of which encode distinct isoforms, including a protease packaged into dense core granules (PC5A) and a type 1 membrane bound protease (PC5B). [provided by RefSeq, May 2014],