



PSA7 rabbit pAb

Catalog_no :	AT7353
Applications :	WB
Reactivity :	Human, Mouse,Rat
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	PSMA7 HSPC
Protein_name :	PSA7
Humangene_id :	<u>5688</u>
Humanswissprot _no :	t <u>O14818</u>
Mousegene_id :	<u>26444</u>
Mouseswissprot _no:	<u>Q9Z2U0</u>
Ratswissprot_no :	<u>P48004</u>
Immunogen :	Synthesized peptide derived from human PSA7
Specificity :	This antibody detects endogenous levels of PSA7 at Human/Mouse/Rat
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Rabbit
Dilution :	WB 1 : 500-2000
Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
Background :	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an

ATP/ubiquitin-dependent process in a non-lysosomal pathway. This gene encodes a member of the peptidase T1A family that functions as a 20S core alpha subunit. The encoded protein interacts with the hepatitis B virus X protein and plays a role in regulating hepatitis C virus internal ribosome entry site (IRES) activity, an activity essential for viral replication. The encoded protein also plays a role in the cellular stress response by regulating hypoxia-inducible factor-1alpha. A pseudogene of this gene is located on the long arm of chromosome 9. [provided by RefSeq, Jul 2012],