



## WFD13 rabbit pAb

Catalog_no :	AT7450
Applications :	IHC-p
Reactivity :	Human, Mouse
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	WFDC13 C20orf138 WAP13
Protein_name :	WFD13
Humangene_id :	<a href="#">164237</a>
Humanswissprot_no :	<a href="#">Q8IUB5</a>
Mousegene_id :	<a href="#">408190</a>
Mouseswissprot_no :	<a href="#">Q5DQQ6</a>
Immunogen :	Synthesized peptide derived from human WFD13
Specificity :	This antibody detects endogenous levels of WFD13 at Human/Mouse
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Rabbit
Dilution :	IHC-p 1 : 50-200
Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
Background :	This gene encodes a member of the WAP-type four-disulfide core (WFDC) domain family. The WFDC domain, or WAP signature motif, contains eight cysteines forming four disulfide bonds at the core of the protein, and functions as a protease inhibitor. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the telomeric cluster. [provided by RefSeq, Jul 2008],



**专注二抗产品**

Biodragon旗下品牌

---