

## DUS7 rabbit pAb

Catalog\_no: AT7554

Applications: WB

Reactivity: Human, Mouse, Rat

Category: 抗原抗体

Size: 100μg/50μg/20μg

Gene\_name: **DUSP7 PYST2** 

Protein\_name: DUS7

Humangene\_id 1849

Humanswissprot Q16829

\_no:

Mousegene\_id: 235584

Mouseswissprot Q91Z46

\_no:

Ratswissprot\_no Q63340

Immunogen: Synthesized peptide derived from human DUS7

Specificity: This antibody detects endogenous levels of DUS7 at Human/Mouse/Rat

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Formulation:

Source: Rabbit

Dilution: WB 1:500-2000

The antibody was affinity-purified from rabbit serum by affinity-chromatography using Purification:

specific immunogen.

Concentration: 1 mg/ml

Storage\_stability -20°C/1 year

Background: Dual-specificity phosphatases (DUSPs) constitute a large heterogeneous subgroup of the

type I cysteine-based protein-tyrosine phosphatase superfamily. DUSPs are

characterized by their ability to dephosphorylate both tyrosine and serine/threonine residues. DUSP7 belongs to a class of DUSPs, designated MKPs, that dephosphorylate MAPK (mitogen-activated protein kinase) proteins ERK (see MIM 601795), JNK (see MIM



601158), and p38 (see MIM 600289) with specificity distinct from that of individual MKP proteins. MKPs contain a highly conserved C-terminal catalytic domain and an N-terminal Cdc25 (see MIM 116947)-like (CH2) domain. MAPK activation cascades mediate various physiologic processes, including cellular proliferation, apoptosis, differentiation, and stress responses (summary by Patterson et al., 2009 [PubMed 19228121]).[supplied by OMIM, Dec 2009],