

FBX17 rabbit pAb

Catalog_no :	<u>AT7637</u>
Applications :	<u>WB</u>
Reactivity :	<u>Human, Mouse,Rat</u>
Category :	<u>抗原抗体</u>
Size :	<u>100µg/50µg/20µg</u>
Gene_name :	<u>FBXO17 FBG4 FBX17 FBX26 FBXO26</u>
Protein_name :	<u>FBX17</u>
Humangene_id :	<u>115290</u>
Humanswissprot_no :	<u>Q96EF6</u>
Mousegene_id :	<u>50760</u>
Mouseswissprot_no :	<u>Q9QZM8</u>
Ratgene_id :	<u>292757</u>
Ratswissprot_no :	<u>Q6AY27</u>
Immunogen :	<u>Synthesized peptide derived from human FBX17</u>
Specificity :	<u>This antibody detects endogenous levels of FBX17 at Human/Mouse/Rat</u>
Formulation :	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
Source :	<u>Rabbit</u>
Dilution :	<u>WB 1 : 500-2000</u>
Purification :	<u>The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage_stability :	<u>-20°C/1 year</u>
Background :	<u>This gene encodes a member of the F-box protein family which is characterized by the F-box motif. The F-box proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in</u>

phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class and it contains an F-box domain. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013],
