

LY65C rabbit pAb

| | |
|---------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Catalog_no : | AT7655 |
| Applications : | WB |
| Reactivity : | Human, Mouse,Rat |
| Category : | 抗原抗体 |
| Size : | 100µg/50µg/20µg |
| Gene_name : | LY6G5C C6orf20 G5C NG33 |
| Protein_name : | LY65C |
| Humangene_id : | 80741 |
| Humanswissprot_no : | Q5SRR4 |
| Mousegene_id : | 114652 |
| Mouseswissprot_no : | Q8K1T5 |
| Ratswissprot_no : | Q8CHN2 |
| Immunogen : | Synthesized peptide derived from human LY65C |
| Specificity : | This antibody detects endogenous levels of LY65C at Human/Mouse/Rat |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Rabbit |
| Dilution : | WB 1 : 500-2000 |
| Purification : | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability : | -20°C/1 year |
| Background : | LY6G5C belongs to a cluster of leukocyte antigen-6 (LY6) genes located in the major histocompatibility complex (MHC) class III region on chromosome 6. Members of the LY6 superfamily typically contain 70 to 80 amino acids, including 8 to 10 cysteines. Most LY6 proteins are attached to the cell surface by a glycosylphosphatidylinositol (GPI) anchor that is directly involved in signal transduction (Mallya et al., 2002 [PubMed]) |



12079290)].[supplied by OMIM, Mar 2008],
