

PCDB7 rabbit pAb

Catalog_no: AT7680

Applications: WB

Reactivity: Human

Category: 抗原抗体

Size: $100 \mu g/50 \mu g/20 \mu g$

Gene_name : PCDHB7

Protein_name : PCDB7

Humangene_id 56129

:

Humanswissprot Q9Y5E2

no:

Immunogen: Synthesized peptide derived from human PCDB7

Specificity: This antibody detects endogenous levels of PCDB7 at Human

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Rabbit

Dilution: WB 1:500-2000

Purification: The antibody was affinity-purified from rabbit serum by affinity-chromatography using

specific immunogen.

Concentration: 1 mg/ml

Storage_stability -20°C/1 year

:

Background: This gene is a member of the protocadherin beta gene cluster, one of three related gene

clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the transcripts from these genes are made up of only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific cell-cell neural connections. The transcript for this particular family member uses more than one polyadenylation site. [provided by RefSeq, Jul 2008],

