

## CSAD rabbit pAb

|                   |   |
|-------------------|---|
| Catalog_no :      | AT7700  |
| Applications :    | WB  |
| Reactivity :      | Human, Mouse,Rat  |
| Category :        | 抗原抗体  |
| Size :            | 100µg/50µg/20µg   |
| Gene_name :       | CSAD CSD  |
| Protein_name :    | CSAD  |
| Humangene_id      | <a href="#">51380</a>   |
| :                 |   |
| Humanswissprot    | <a href="#">Q9Y600</a>  |
| _no :             |   |
| Mousegene_id :    | <a href="#">246277</a>  |
| Mouseswissprot    | <a href="#">Q9DBE0</a>  |
| _no :             |   |
| Ratgene_id :      | <a href="#">60356</a>   |
| Ratswissprot_no   | <a href="#">Q64611</a>  |
| :                 |   |
| Immunogen :       | Synthesized peptide derived from human CSAD   |
| Specificity :     | This antibody detects endogenous levels of CSAD at Human/Mouse/Rat  |
| Formulation :     | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source :          | Rabbit  |
| Dilution :        | WB 1 : 500-2000   |
| Purification :    | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.   |
| Concentration :   | 1 mg/ml   |
| Storage_stability | -20°C/1 year  |
| :                 |   |
| Background :      | This gene encodes a member of the group 2 decarboxylase family. A similar protein in rodents plays a role in multiple biological processes as the rate-limiting enzyme in taurine biosynthesis, catalyzing the decarboxylation of cysteinesulfinate to hypotaurine. |



Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Sep 2011],

---