



## SRC-3 (phospho-Thr24) rabbit pAb

Catalog_no :	AP1510
Applications :	WB
Reactivity :	Human
Category :	抗原抗体
Size :	100µg/50µg/20µg
Gene_name :	NCOA3 AIB1 BHLHE42 RAC3 TRAM1
Protein_name :	SRC-3 (Thr24)
Humangene_id :	<u>8202</u>
Humanswissprot _no :	2 <u>Q9Y6Q9</u>
Mouseswissprot _no :	<u>009000</u>
Ratswissprot_no :	<u>Q9EPU2</u>
Immunogen :	Synthesized phosho peptide around human SRC-3 (Thr24)
Specificity :	This antibody detects endogenous levels of Human SRC-3 (phospho-Thr24)
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Rabbit
Dilution :	WB 1:1000-2000
Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Concentration :	1 mg/ml
Storage_stability :	-20°C/1 year
Other_name :	Nuclear receptor coactivator 3 (NCoA-3) (EC 2.3.1.48) (ACTR) (Amplified in breast cancer 1 protein) (AIB-1) (CBP-interacting protein) (pCIP) (Class E basic helix-loop-helix protein 42) (bHLHe42) (Receptor-associated coactivator 3) (RAC-3) (Steroid receptor coactivator protein 3) (SRC-3) (Thyroid hormone receptor activator molecule 1) (TRAM-1)
Molecular Weight :	160KD



Background : nuclear receptor coactivator 3(NCOA3) Homo sapiens The protein encoded by this gene is a nuclear receptor coactivator that interacts with nuclear hormone receptors to enhance their transcriptional activator functions. The encoded protein has histone acetyltransferase activity and recruits p300/CBP-associated factor and CREB binding protein as part of a multisubunit coactivation complex. This protein is initially found in the cytoplasm but is translocated into the nucleus upon phosphorylation. Several transcript variants encoding different isoforms have been found for this gene. In addition, a polymorphic repeat region is found in the C-terminus of the encoded protein. [provided by RefSeq, Mar 2010],